

THE NORM OF A TRUNCATED TOEPLITZ OPERATOR

STEPHAN RAMON GARCIA AND WILLIAM T. ROSS

ABSTRACT. We prove several lower bounds for the norm of a truncated Toeplitz operator and obtain a curious relationship between the H^2 and H^∞ norms of functions in model spaces.

1. INTRODUCTION

In this paper, we continue the discussion initiated in [6] concerning the norm of a truncated Toeplitz operator. In the following, let H^2 denote the classical Hardy space of the open unit disk $\mathbb{D}$ and $K_\Theta := H^2 \cap (\Theta H^2)^\perp$, where Θ is an inner function, denote one of the so-called Jordan model spaces [2, 4, 7]. If H^∞ is the set of all bounded analytic functions on $\mathbb{D}$, the space $K_\Theta^\infty := H^\infty \cap K_\Theta$ is norm dense in K_Θ (see [2, p. 83] or [9, Lem. 2.3]). If P_Θ is the orthogonal projection from $L^2 := L^2(\partial\mathbb{D}, \frac{|d\zeta|}{2\pi})$ onto K_Θ and $\varphi \in L^2$, then the operator

$$A_\varphi f := P_\Theta(\varphi f), \quad f \in K_\Theta^\infty,$$

is densely defined on K_Θ and is called a *truncated Toeplitz operator*. Various aspects of these operators were studied in [3, 5, 6, 9, 10].

If $\|\cdot\|$ is the norm on L^2 , we let

$$\|A_\varphi\| := \sup\{\|A_\varphi f\| : f \in K_\Theta^\infty, \|f\| = 1\} \quad (1)$$

and note that this quantity is finite if and only if A_φ extends to a bounded operator on K_Θ . When $\varphi \in L^\infty$, the set of bounded measurable functions on $\partial\mathbb{D}$, we have the basic estimates

$$0 \leq \|A_\varphi\| \leq \|\varphi\|_\infty.$$

However, it is known that equality can hold, in nontrivial ways, in either of the inequalities above and hence finding the norm of a truncated Toeplitz operator can be difficult. Furthermore, it turns out that there are many unbounded symbols $\varphi \in L^2$ which yield bounded operators A_φ . Unlike the situation for classical Toeplitz operators on H^2 , for a given $\varphi \in L^2$, there many $\psi \in L^2$ for which $A_\varphi = A_\psi$ [9, Thm. 3.1].

For a given symbol $\varphi \in L^2$ and inner function Θ , lower bounds on the quantity (1) are useful in answering the following nontrivial questions:

- (i) is A_φ unbounded?
- (ii) if $\varphi \in L^\infty$, is A_φ norm-attaining (i.e., is $\|A_\varphi\| = \|\varphi\|_\infty$)?

2000 *Mathematics Subject Classification.* 47A05, 47B35, 47B99.

Key words and phrases. Toeplitz operator, model space, truncated Toeplitz operator, reproducing kernel, complex symmetric operator, conjugation.

First author partially supported by National Science Foundation Grant DMS-0638789.

When Θ is a finite Blaschke product and $\varphi \in H^\infty$, the paper [6] computes $\|A_\varphi\|$ and gives necessary and sufficient conditions as to when $\|A_\varphi\| = \|\varphi\|_\infty$. The purpose of this short note is to give a few lower bounds on $\|A_\varphi\|$ for general inner functions Θ and general $\varphi \in L^2$. Along the way, we obtain a curious relationship (Corollary 5) between the H^2 and H^∞ norms of functions in K_Θ^∞ .

2. LOWER BOUNDS DERIVED FROM POISSON'S FORMULA

For $\varphi \in L^2$, let

$$(\mathfrak{P}\varphi)(z) := \int_{\partial\mathbb{D}} \frac{1 - |z|^2}{|\zeta - z|^2} \varphi(\zeta) \frac{|d\zeta|}{2\pi}, \quad z \in \mathbb{D}, \quad (2)$$

be the standard Poisson extension of φ to $\mathbb{D}$. For $\varphi \in C(\partial\mathbb{D})$, the continuous functions on $\partial\mathbb{D}$, recall that $\mathfrak{P}\varphi$ solves the classical Dirichlet problem with boundary data φ . Also note that

$$k_\lambda(z) := \frac{1 - \overline{\Theta(\lambda)}\Theta(z)}{1 - \overline{\lambda}z}, \quad \lambda, z \in \mathbb{D},$$

is the reproducing kernel for K_Θ [9].

Our first result provides a general lower bound for $\|A_\varphi\|$ which yields a number of useful corollaries:

Theorem 1. *If $\varphi \in L^2$, then*

$$\sup_{\lambda \in \mathbb{D}} \frac{1 - |\lambda|^2}{1 - |\Theta(\lambda)|^2} \left| \int_{\partial\mathbb{D}} \varphi(z) \left| \frac{\Theta(z) - \Theta(\lambda)}{z - \lambda} \right|^2 \frac{|dz|}{2\pi} \right| \leq \|A_\varphi\|. \quad (3)$$

In other words,

$$\sup_{\lambda \in \mathbb{D}} \left| \int_{\partial\mathbb{D}} \varphi(z) d\nu_\lambda(z) \right| \leq \|A_\varphi\|$$

where

$$d\nu_\lambda(z) := \frac{1 - |\lambda|^2}{1 - |\Theta(\lambda)|^2} \left| \frac{\Theta(z) - \Theta(\lambda)}{z - \lambda} \right|^2 \frac{|dz|}{2\pi}$$

is a family of probability measures on $\partial\mathbb{D}$ indexed by $\lambda \in \mathbb{D}$.

Proof. For $\lambda \in \mathbb{D}$ we have

$$\|k_\lambda\| = \sqrt{\frac{1 - |\Theta(\lambda)|^2}{1 - |\lambda|^2}}, \quad (4)$$

from which it follows that

$$\begin{aligned} \|A_\varphi\| &\geq \frac{1 - |\lambda|^2}{1 - |\Theta(\lambda)|^2} |\langle A_\varphi k_\lambda, k_\lambda \rangle| \\ &= \frac{1 - |\lambda|^2}{1 - |\Theta(\lambda)|^2} |\langle P_\Theta \varphi k_\lambda, k_\lambda \rangle| \\ &= \frac{1 - |\lambda|^2}{1 - |\Theta(\lambda)|^2} |\langle \varphi k_\lambda, k_\lambda \rangle| \\ &= \frac{1 - |\lambda|^2}{1 - |\Theta(\lambda)|^2} \left| \int_{\partial\mathbb{D}} \varphi(z) \left| \frac{\Theta(z) - \Theta(\lambda)}{z - \lambda} \right|^2 \frac{|dz|}{2\pi} \right|. \end{aligned}$$

That the measures $d\nu_\lambda$ are indeed probability measures follows from (4). $\square$

Now observe that if $\Theta(\lambda) = 0$, the argument in the supremum on the left hand side of (3) becomes the absolute value of the expression in (2). This immediately yields the following corollary:

Corollary 1. *If $\varphi \in L^2$, then*

$$\sup_{\lambda \in \Theta^{-1}(\{0\})} |(\mathfrak{P}\varphi)(\lambda)| \leq \|A_\varphi\|, \quad (5)$$

where the supremum is to be regarded as 0 if $\Theta^{-1}(\{0\}) = \emptyset$.

Under the right circumstances, the preceding corollary can be used to prove that certain truncated Toeplitz operators are norm-attaining:

Corollary 2. *Let Θ be an inner function having zeros which accumulate at every point of $\partial\mathbb{D}$. If $\varphi \in C(\partial\mathbb{D})$ then $\|A_\varphi\| = \|\varphi\|_\infty$.*

Proof. Let $\zeta \in \partial\mathbb{D}$ be such that $|\varphi(\zeta)| = \|\varphi\|_\infty$. By hypothesis, there exists a sequence λ_n of zeros of Θ which converge to ζ . By continuity, we conclude that

$$\|\varphi\|_\infty = \lim_{n \rightarrow \infty} |(\mathfrak{P}\varphi)(\lambda_n)| \leq \|A_\varphi\| \leq \|\varphi\|_\infty$$

whence $\|A_\varphi\| = \|\varphi\|_\infty$. $\square$

The preceding corollary stands in contrast to the finite Blaschke product setting. Indeed, if Θ is a finite Blaschke product and $\varphi \in H^\infty$, then it is known that $\|A_\varphi\| = \|\varphi\|_\infty$ if and only if φ is the scalar multiple of the inner factor of some function from K_Θ [6, Thm. 2].

At the expense of wordiness, the hypothesis of Corollary 2 can be considerably weakened. A cursory examination of the proof indicates that we only need ζ to be a limit point of the zeros of Θ , $\varphi \in L^\infty$ to be continuous on an open arc containing ζ , and $|\varphi(\zeta)| = \|\varphi\|_\infty$.

Theorem 1 yields yet another lower bound for $\|A_\varphi\|$. Recall that an inner function Θ has a finite angular derivative at $\zeta \in \partial\mathbb{D}$ if Θ has a non-tangential limit $\Theta(\zeta)$ of modulus one at ζ and Θ' has a finite non-tangential limit $\Theta'(\zeta)$ at ζ . This is equivalent to asserting that

$$\frac{\Theta(z) - \Theta(\zeta)}{z - \zeta} \quad (6)$$

has the non-tangential limit $\Theta'(\zeta)$ at ζ . If Θ has a finite angular derivative at ζ , then the function in (6) belongs to H^2 and

$$\lim_{r \rightarrow 1^-} \int_{\partial\mathbb{D}} \left| \frac{\Theta(z) - \Theta(r\zeta)}{z - r\zeta} \right|^2 \frac{|dz|}{2\pi} = \int_{\partial\mathbb{D}} \left| \frac{\Theta(z) - \Theta(\zeta)}{z - \zeta} \right|^2 \frac{|dz|}{2\pi}.$$

Furthermore, the above is equal to

$$\lim_{r \rightarrow 1^-} \frac{1 - |\Theta(r\zeta)|^2}{1 - r^2} = |\Theta'(\zeta)| > 0.$$

See [1, 8] for further details on angular derivatives. Theorem 1 along with the preceding discussion and Fatou's lemma yield the following lower estimate for $\|A_\varphi\|$.

Corollary 3. *For an inner function Θ , let D_Θ be the set of $\zeta \in \partial\mathbb{D}$ for which Θ has a finite angular derivative $\Theta'(\zeta)$ at ζ . If $\varphi \in L^\infty$ or if $\varphi \in L^2$ with $\varphi \geq 0$, then*

$$\sup_{\zeta \in D_\Theta} \frac{1}{|\Theta'(\zeta)|} \left| \int_{\partial\mathbb{D}} \varphi(z) \left| \frac{\Theta(z) - \Theta(\zeta)}{z - \zeta} \right|^2 \frac{|dz|}{2\pi} \right| \leq \|A_\varphi\|.$$

In other words,

$$\sup_{\zeta \in D_\Theta} \left| \int_{\partial\mathbb{D}} \varphi(z) d\nu_\lambda(z) \right| \leq \|A_\varphi\|,$$

where

$$d\nu_\lambda(z) := \frac{1}{|\Theta'(\zeta)|} \left| \frac{\Theta(z) - \Theta(\zeta)}{z - \zeta} \right|^2 \frac{|dz|}{2\pi}$$

is a family of probability measures on $\partial\mathbb{D}$ indexed by $\zeta \in D_\Theta$.

3. LOWER BOUNDS AND PROJECTIONS

Our next several results concern lower bounds on $\|A_\varphi\|$ involving the orthogonal projection $P_\Theta : L^2 \rightarrow K_\Theta$.

Theorem 2. *If Θ is an inner function and $\varphi \in L^2$, then*

$$\frac{\|P_\Theta(\varphi) - \overline{\Theta(0)}P_\Theta(\Theta\varphi)\|}{(1 - |\Theta(0)|^2)^{\frac{1}{2}}} \leq \|A_\varphi\|.$$

Proof. First observe that $\|k_0\| = (1 - |\Theta(0)|^2)^{\frac{1}{2}}$. Next we see that if $\varphi \in L^2$ and $g \in K_\Theta$ is any unit vector, then

$$\begin{aligned} (1 - |\Theta(0)|^2)^{\frac{1}{2}} \|A_\varphi\| &\geq |\langle A_\varphi k_0, g \rangle| \\ &= |\langle P_\Theta(\varphi k_0), g \rangle| \\ &= |\langle P_\Theta(\varphi) - \overline{\Theta(0)}P_\Theta(\Theta\varphi), g \rangle|. \end{aligned}$$

Setting

$$g = \frac{P_\Theta(\varphi) - \overline{\Theta(0)}P_\Theta(\Theta\varphi)}{\|P_\Theta(\varphi) - \overline{\Theta(0)}P_\Theta(\Theta\varphi)\|}$$

yields the desired inequality. $\square$

In light of the fact that $P_\Theta(\Theta\varphi) = 0$ whenever $\varphi \in H^2$, Theorem 2 leads us immediately to the following corollary:

Corollary 4. *If Θ is inner and $\varphi \in H^2$, then*

$$\frac{\|P_\Theta(\varphi)\|}{(1 - |\Theta(0)|^2)^{1/2}} \leq \|A_\varphi\|. \quad (7)$$

It turns out that (7) has a rather interesting function-theoretic implication. Let us first note that for $\varphi \in H^\infty$, we can expect no better inequality than

$$\|\varphi\| \leq \|\varphi\|_\infty$$

(with equality holding if and only if φ is a scalar multiple of an inner function). However, if φ belongs to K_Θ^∞ , then a stronger inequality holds.

Corollary 5. *If Θ is an inner function, then*

$$\|\varphi\| \leq (1 - |\Theta(0)|^2)^{\frac{1}{2}} \|\varphi\|_\infty \quad (8)$$

holds for all $\varphi \in K_\Theta^\infty$. If Θ is a finite Blaschke product, then equality holds if and only if φ is a scalar multiple of an inner function from K_Θ .

Proof. First observe that the inequality

$$\|\varphi\| \leq (1 - |\Theta(0)|^2)^{\frac{1}{2}} \|\varphi\|_\infty$$

follows from Corollary 4 and the fact that $P_\Theta \varphi = \varphi$ whenever $\varphi \in K_\Theta$. Now suppose that Θ is a finite Blaschke product and assume that equality holds in the preceding for some $\varphi \in K_\Theta^\infty$. In light of (7), it follows that $\|A_\varphi\| = \|\varphi\|_\infty$. From [6, Thm. 2] we see that φ must be a scalar multiple of the inner *part* of a function from K_Θ . But since $\varphi \in K_\Theta^\infty$, then φ must be a scalar multiple of an inner function from K_Θ . $\square$

When Θ is a finite Blaschke product, then K_Θ is a finite dimensional subspace of H^2 consisting of bounded functions [3, 5, 9]. By elementary functional analysis, there are $c_1, c_2 > 0$ so that

$$c_1 \|\varphi\| \leq \|\varphi\|_\infty \leq c_2 \|\varphi\|$$

for all $\varphi \in K_\Theta$. This prompts the following question:

Question. What are the optimal constants c_1, c_2 in the above inequality?

4. LOWER BOUNDS FROM THE DECOMPOSITION OF K_Θ

A result of Sarason [9, Thm. 3.1] says, for $\varphi \in L^2$, that

$$A_\varphi \equiv 0 \Leftrightarrow \varphi \in \Theta H^2 + \overline{\Theta H^2}. \quad (9)$$

It follows that the most general truncated Toeplitz operator on K_Θ is of the form $A_{\psi+\bar{\chi}}$ where $\psi, \chi \in K_\Theta$. We can refine this observation a bit further and provide another canonical decomposition for the symbol of a truncated Toeplitz operator.

Lemma 1. *Each bounded truncated Toeplitz operator on K_Θ is generated by a symbol of the form*

$$\varphi = \underbrace{\psi}_{\in H^2} + \underbrace{\chi \bar{\Theta}}_{\in zH^2} \quad (10)$$

where $\psi, \chi \in K_\Theta$.

Before getting to the proof, we should remind the reader of a technical detail. It follows from the identity $K_\Theta = H^2 \cap \Theta \overline{zH^2}$ (see [2, p. 82]) that

$$C : K_\Theta \rightarrow K_\Theta, \quad Cf := \overline{zf} \Theta,$$

is an isometric, conjugate-linear, involution. In fact, when A_φ is a bounded operator we have the identity $CA_\varphi C = A_\varphi^*$ [9, Lemma 2.1].

Proof of Lemma 1. If T is a bounded truncated Toeplitz operator on K_Θ , then there exists some $\varphi \in L^2$ such that $T = A_\varphi$. We claim that this φ can be chosen to have the special form (10). First let us write $\varphi = f + \overline{zg}$ where $f, g \in H^2$. Using the orthogonal decomposition $H^2 = K_\Theta \oplus \Theta H^2$, it follows that φ may be further decomposed as

$$\varphi = (f_1 + \Theta f_2) + \overline{z(g_1 + \Theta g_2)}$$

where $f_1, g_1 \in K_\Theta$ and $f_2, g_2 \in H^2$. By (9), the symbols Θf_2 and $\overline{\Theta(zg_2)}$ yield the zero truncated Toeplitz operator on K_Θ . Therefore we may assume that

$$\varphi = f + \overline{zg}$$

for some $f, g \in K_\Theta$. Since $Cg = \overline{gz} \Theta$, we have $\overline{zg} = (Cg) \bar{\Theta}$ and hence (10) holds with $\psi = f$ and $\chi = Cg$. $\square$

Corollary 6. *Let Θ be an inner function. If $\psi_1, \psi_2 \in K_\Theta$ and $\varphi = \psi_1 + \psi_2 \overline{\Theta}$, then*

$$\frac{\|\psi_1 - \overline{\Theta(0)}\psi_2\|}{(1 - |\Theta(0)|^2)^{\frac{1}{2}}} \leq \|A_\varphi\|.$$

Proof. If $\varphi = \psi_1 + \psi_2 \overline{\Theta}$, then, since $\psi_1, \psi_2 \in K_\Theta$ and $\psi_2 \overline{\Theta} \in \overline{zH^2}$, we have

$$P_\Theta(\varphi) - \overline{\Theta(0)}P_\Theta(\Theta\varphi) = \psi_1 - \overline{\Theta(0)}\psi_2.$$

The result now follows from Theorem 2. □

REFERENCES

1. J. A. Cima, A. L. Matheson, and W. T. Ross, *The Cauchy transform*, Mathematical Surveys and Monographs, vol. 125, American Mathematical Society, Providence, RI, 2006.
2. J. A. Cima and W. T. Ross, *The backward shift on the Hardy space*, Mathematical Surveys and Monographs, vol. 79, American Mathematical Society, Providence, RI, 2000.
3. J. A. Cima, W. T. Ross, and W. R. Wogen, *Truncated Toeplitz operators on finite dimensional spaces*, Oper. Matrices **2** (2008), no. 3, 357–369.
4. R. G. Douglas, H. S. Shapiro, and A. L. Shields, *Cyclic vectors and invariant subspaces for the backward shift operator.*, Ann. Inst. Fourier (Grenoble) **20** (1970), no. fasc. 1, 37–76.
5. S. R. Garcia, *Conjugation and Clark operators*, Recent advances in operator-related function theory, Contemp. Math., vol. 393, Amer. Math. Soc., Providence, RI, 2006, pp. 67–111.
6. S. R. Garcia and W. T. Ross, *A nonlinear extremal problem on the Hardy space*, Comp. Methods. Function Theory **9** (2009), no. 2, 485–524.
7. N. Nikolski, *Treatise on the shift operator*, Springer-Verlag, Berlin, 1986.
8. D. Sarason, *Sub-Hardy Hilbert spaces in the unit disk*, University of Arkansas Lecture Notes in the Mathematical Sciences, 10, John Wiley & Sons Inc., New York, 1994, A Wiley-Interscience Publication.
9. ———, *Algebraic properties of truncated Toeplitz operators*, Operators and Matrices **1** (2007), no. 4, 491–526.
10. ———, *Unbounded Toeplitz operators*, Integral Equations Operator Theory **61** (2008), no. 2, 281–298.

DEPARTMENT OF MATHEMATICS, POMONA COLLEGE, CLAREMONT, CALIFORNIA, 91711, USA
E-mail address: `Stephan.Garcia@pomona.edu`
URL: `http://pages.pomona.edu/~sg064747`

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, UNIVERSITY OF RICHMOND, RICHMOND, VIRGINIA, 23173, USA
E-mail address: `wross@richmond.edu`
URL: `http://facultystaff.richmond.edu/~wross`